

**LINDNER
610 EARLY WEAN 22 MX BN
MEDICATED**

A complete feed for Starting Pigs being fed for Exhibition. For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections of swine.

ACTIVE DRUG INGREDIENT

Carbadox0.0055% (50 grams per ton)
 Pyrantel Tartrate.....0.0106% (96 grams per ton)

GUARANTEED ANALYSIS

Crude Protein – Min..... 22.0%
 Lysine—Min.....1.73%
 Crude Fat –Min..... 5.6%
 Crude Fiber—Max..... 2.4%
 Calcium—Min..... 0.8%
 Calcium—Max..... 1.0%
 Phosphorous—Min..... 0.64%
 Salt—Min.....None added
 Selenium—Min.....0.3 ppm
 Zinc—Min 3700 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Early wean 22% is a complete feed specifically formulated for 8 to 15 lb. pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 15lb body weight. Then they should be fed an approved starter ration.

WARNING

Do not feed to swine weighing more than 75 lb. body weight. Do not feed to swine within 10 weeks of slaughter.

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

MANUFACTURED BY
LINDNER FEED & MILLING CO., INC.
 Comfort, TX 78013
 NET WT 50 LB (22.68KG)

**LINDNER
611 Scud Missile
MEDICATED**

A complete feed for Starting Pigs being fed for Exhibition. For increased rate of weight gain and improved feed efficiency.

ACTIVE DRUG INGREDIENT

Chlortetracycline 50 Grams per ton

GUARANTEED ANALYSIS

Crude Protein – Min..... 23.0%
 Lysine—Min.....1.42%
 Crude Fat –Min..... 7.10%
 Crude Fiber—Max..... 4.20%
 Calcium—Min..... 0.78%
 Calcium—Max..... 0.85%
 Phosphorous—Min..... 0.67%
 Salt—Min.....0.01%
 Selenium—Min.....0.3 ppm
 Zinc—Min 3200 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Soybean Hulls, Dried Whey, Yeast Culture, Plasma, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

611 Scud Missile is a complete feed specifically formulated for 8 to 15 lb. pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 15lb body weight. Then they should be fed an approved starter ration.

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

MANUFACTURED BY
LINDNER FEED & MILLING CO., INC.
 Comfort, TX 78013
 NET WT 50 LB (22.68KG)

**LINDNER
612 Scud Missile
MEDICATED**

A complete feed for Starting Pigs being fed for Exhibition. For the control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin. Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENT

Tiamulin hydrogen fumarate 35 g/ton
 Chlortetracycline 400 g/ton

GUARANTEED ANALYSIS

Crude Protein – Min..... 23.0%
 Lysine—Min.....1.42%
 Crude Fat –Min..... 7.10%
 Crude Fiber—Max..... 4.20%
 Calcium—Min..... 0.78%
 Calcium—Max..... 0.85%
 Phosphorous—Min..... 0.67%
 Salt—Min.....0.01%
 Selenium—Min.....0.3 ppm
 Zinc—Min 3200 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Soybean Hulls, Dried Whey, Yeast Culture, Plasma, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

612 Scud Missile is a complete feed specifically formulated for 8 to 15 lb. pigs fed for exhibition. This product must be fed continuously as the sole ration for 14 days only. Then they should be fed an approved starter ration.

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite. Do not feed to pigs while they are on any other medicated feed. If signs of toxicity occur, discontinue use of medicated feeds. Not for use in swine weighing more than 250 lb.

MANUFACTURED BY
LINDNER FEED & MILLING CO., INC.
 Comfort, TX 78013
 NET WT 50 LB (22.68KG)

LINDNER

615 Pig Starter 20% MX BN

MEDICATED

A complete feed for Starting Pigs being fed for Exhibition. For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections of swine.

ACTIVE DRUG INGREDIENT

Carbadox0.0055% (50 grams per ton)
Pyrantel Tartrate.....0.0106% (96 grams per ton)

GUARANTEED ANALYSIS

Crude Protein – Min.....20.0%
Lysine—Min.....1.63%
Crude Fat –Min.....5.6%
Crude Fiber—Max.....2.6%
Calcium—Min.....1.0%
Calcium—Max.....1.1%
Phosphorous—Min.....0.64%
Salt—Min.....None added
Selenium—Min.....0.3 ppm
Zinc—Min.....2100 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Pig Starter 20% is a complete feed specifically formulated for pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 50 lb. body weight. Then they should be fed an approved grower ration.

WARNING

Do not feed to swine weighing more than 75 lb. body weight. Do not feed to swine within 10 weeks of slaughter.

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

MANUFACTURED BY
LINDNER FEED & MILLING CO., INC.
Comfort, TX 78013
NET WT 50 LB (22.68KG)

LINDNER

630 Pig Starter Grower MX/BN

MEDICATED

A complete feed for Starting Pigs being fed for Exhibition. For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections of swine.

ACTIVE DRUG INGREDIENT

Carbadox0.0055% (50 grams per ton)
Pyrantel Tartrate.....0.0106% (96 grams per ton)

GUARANTEED ANALYSIS

Crude Protein – Min.....20.0%
Lysine—Min.....1.3%
Crude Fat –Min.....4.6%
Crude Fiber—Max.....2.7%
Calcium—Min.....0.6%
Calcium—Max.....0.7%
Phosphorous—Min.....0.5%
Salt—Min.....0.2%
Selenium—Min.....0.3 ppm
Zinc—Min.....160 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Pig Starter Grower is a complete feed specifically formulated for pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 75-lb body weight. Then they should be fed an approved grower finisher ration.

WARNING

Do not feed to swine weighing more than 75 lb body weight. Do not feed to swine within 10 weeks of slaughter.

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

MANUFACTURED BY
LINDNER FEED & MILLING CO., INC.
Comfort, TX 78013
NET WT 50 LB (22.68kg)

LINDNER

631 Starter / Grower TY

MEDICATED

A complete feed for Starter /Grower Pigs being fed for Exhibition. Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis.

ACTIVE DRUG INGREDIENT

Tylosin.....(50 mg/lb.).....100 gm/ton

GUARANTEED ANALYSIS

Crude Protein – Min.....20.0%
Lysine—Min.....1.3%
Crude Fat –Min.....4.6%
Crude Fiber—Max.....2.7%
Calcium—Min.....0.6%
Calcium—Max.....0.7%
Phosphorous—Min.....0.5%
Salt—Min.....0.2%
Salt—Max.....0.3%
Selenium—Min.....0.3 ppm
Zinc—Min.....160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Lindner's Starter Grower TY is a complete feed specifically formulated for pigs fed for exhibition. It should be fed continuously as the sole Medication to pigs weighing not less than 45 lb. body weight. Specific feeding instructions may vary according to the goals of the feeding program for show pigs. Generally, pigs can be fed ad lib, or feed may be restricted to achieve desired growth rates.

MANUFACTURED BY
LINDNER FEED AND MILLING CO. INC.

Comfort, Texas 78013
NET WT.50 LB (22.68 kg)

LINDNER
632 Starter / Grower LI-FBZ
MEDICATED

A complete feed for Starter/Grower Pigs being fed for Exhibition. For reduction in the severity of swine Mycoplasma pneumonia caused by *Mycoplasma hyopneumoniae*. For the removal of: Lung worms: *Metastrongylus apri*, *Metastrongylus pudendotectus*. Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); nodular (*Oesophagostomum dentatum*, *Ornithostrongylus quadrispinulatum*); small stomach (*Hyostrogylus rubidus*); Adult and larvae (L2, L3, L4 stages intestinal mucosal forms) whipworm (*Trichuris suis*); Kidneyworm: Adult and larvae *Stephanurus dentatus*.

ACTIVE DRUG INGREDIENT

Lincomycin 200 gm/ton
 Fenbendazole 29.2 gm/ton (14.6 mg/lb.)

GUARANTEED ANALYSIS

Crude Protein – Min.....20.0%
 Lysine—Min.....1.3%
 Crude Fat –Min.....4.6%
 Crude Fiber—Max.....2.7%
 Calcium—Min.....0.6%
 Calcium—Max.....0.7%
 Phosphorous—Min.....0.5%
 Salt—Min.....0.2%
 Salt—Max.....0.3%
 Selenium—Min.....0.3 ppm
 Zinc Min. 160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Feed as the sole ration to provide a total intake of 9mg Fenbendazole/kg of body weight (4.08mg Fenbendazole/lb. of body weight) divided over 3-12 days. Multiply 0.28 times pound of body weight and feed this amount of product over 3-12 days.

Example:

Body weight of Pig to be Dewormed	Starter/Grower Feed Required per pig	Days Required to Consume Dose
50 lb.	14.0 lb.	7
75 lb.	21.0 lb.	7
100 lb.	28.0 lb.	7

CAUTION STATEMENT

Occasionally, swine fed Lincomycin may, within the first 2 days after the onset of treatment, develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5-8 days without discontinuing the Lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feeds containing Lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not to be fed to swine that weigh more than 250 lbs. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasites.

WARNING

Do not slaughter swine for human consumption for 6 days following last treatment with Lincomycin.

MANUFACTURED BY
LINDNER FEED AND MILLING CO. INC.
 Comfort, Texas 78013
 NET WT.50 LB (22.68 kg)

LINDNER
633 Starter / Grower ASP
MEDICATED

A complete feed for Starter / Grower Pigs being fed for Exhibition. Administer to swine in a complete feed for reduction of the incidence of cervical abscesses; treatment of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery); maintenance of weight gains in the presence of atrophic rhinitis; increased rate of weight gain and improved feed efficiency from 10 lb. bodyweight up to 6 weeks postweaning. For swine raised in confinement (dry lot) or on limited pasture.

ACTIVE DRUG INGREDIENT

Chlortetracycline 100 gm/ton
 Sulfathiazole 100 gm/ton
 Penicillin 50 gm/ton

GUARANTEED ANALYSIS

Crude Protein – Min.....20.0%
 Lysine—Min.....1.3%
 Crude Fat –Min.....4.6%
 Crude Fiber—Max.....2.7%
 Calcium—Min.....0.6%
 Calcium—Max.....0.7%
 Phosphorous—Min.....0.5%
 Salt—Min.....0.2%
 Salt—Max.....0.3%
 Selenium—Min.....0.3 ppm
 Zinc—Min..... 160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Lindner's Starter Grower ASP is a complete feed specifically formulated for pigs fed for exhibition. It should be fed continuously as the sole medicated ration to pigs weighing up to 75 lb. body weight. Specific feeding instructions may vary according to the goals of the feeding program for show pigs. The animals should be self fed.

WARNING STATEMENT

Withdraw 7 days prior to slaughter.

MANUFACTURED BY
LINDNER FEED AND MILLING CO. INC.
 Comfort, Texas 78013
 NET WT.50 LB (22.68 kg)

LINDNER
670 Grower TY
MEDICATED

A Complete Feed for Grower Pigs Being Fed for Exhibition.

For maintaining weight gains and feed efficiency in the presence of atrophic rhinitis.

ACTIVE DRUG INGREDIENT

Tylosin.....(50 mg/lb.).....100 gm /ton

GUARENTEED ANALYSIS

Crude Protein—Min.....18.0%
 Lysine—Min..... 1.10%
 Crude Fat –Min..... 4.1%
 Crude Fiber –Max.....2.2%
 Calcium –Min..... 0.5%
 Calcium –Max..... 0.6%
 Phosphorous –Min..... 0.4%
 Salt –Min..... 0.2%
 Salt –Max..... 0.3%
 Selenium—Min..... 0.3 ppm
 Zinc –Min..... 160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Lindner's Grower TY is a complete feed specifically formulated for pigs fed for exhibition. It should be fed Continuously as the sole Medication to pigs weighing 70 lb

body weight to show or finish. Specific feeding instructions may vary according to the goals of the feeding program for show pigs. Generally, pigs can be fed ad lib, or feed may be restricted to achieve desired growth rates.

MANUFACTURED BY

LINDNER FEED AND MILLING CO. INC.
 Comfort, Texas 78013

NET WT. 50 LB (22.68 kg)

**LINDNER
671 Grower BMD-FBZ
MEDICATED**

A complete Feed for Growing Pigs Being Fed for Exhibition

For control of swine dysentery associated with *Brachyspira hyodysenteriae* on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.

FOR THE REMOVAL OF: Lung worms: *Metastrongylus apri*, *Metastrongylus pudendotectus*. Gastrointestinal worms: Adult and larvae (L3,L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); nodular (*Oesophagostomum dentatum*, *Ornithostrongylus quadrispinulatum*); small stomach (*Hyostrongylus rubidus*); Adult and larvae (L2,L3,L4 stages-intestinal mucosal forms) whipworm (*Trichuris suis*). Kidney worm: Adult and larvae *Stephanurus dentatus*.

ACTIVE DRUG INGREDIENTS

Bacitracin Methylene Disalicylate..... 250 g/ton
Fenbendazole..... 29.2 gm/ton(14.6 mg/lb.)

GUARANTEED ANALYSIS

Crude Protein (Min)..... 18.0%
Lysine (Min)..... 1.10%
Crude Fat (Min)..... 4.6%
Crude Fiber (Min)..... 2.3%
Calcium (Min)..... 0.6%
Calcium (Max)..... 0.7%
Phosphorous (Min)..... 0.5%
Salt (Min)..... 0.2%
Salt (Max)..... 0.3%
Selenium (Min)..... 0.3 ppm
Zinc (Min)..... 160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

**SEE BACK OF TAG FOR FEEDING DIRECTIONS,
WARNING AND CAUTION STATEMENTS**

Manufactured By
LINDNER FEED AND MILLING CO. INC.
Comfort, TX 78013
NET WT. 50 LB (22.68 kg)

671 FEEDING DIRECTIONS

Feed as sole ration in complete feed for a period of 3-12 days so to provide a total intake of 9 mg Fenbendazole/kg of body weight. No withdrawal time needed.

AMOUNT NEEDED PER PIG

Feed at the rate of 1 lb. complete feed for each 25 lb. body wt. each day for 7 days.

CAUTION

Not for use in swine that weigh more than 250 lb. Under conditions of continued exposure to parasites, re-treatment may be needed after 4-6 weeks. Diagnosis of swine dysentery should be confirmed by a veterinarian when results are not satisfactory.

**LINDNER
674 Pig Finisher Payleane
MEDICATED**

A Complete Feed for Finisher Pigs Being Fed for Exhibition.

For improved feed efficiency and increased carcass leanness, and for the prevention and/or control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis*.

ACTIVE DRUG INGREDIENT

Ractopamine HCl 18.0 g/ton
Tylosin.....(50 mg/lb).....100 gm /ton

GUARENTEED ANALYSIS

Crude Protein—Min.....18.0%
Lysine—Min..... 1.10%
Crude Fat —Min..... 4.1%
Crude Fiber —Max.....2.2%
Calcium —Min..... 0.5%
Calcium —Max..... 0.6%
Phosphorous —Min..... 0.4%
Salt —Min..... 0.2%
Salt —Max..... 0.3%
Selenium—Min..... 0.3 ppm
Zinc —Min..... 160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Feed continuously to finishing swine as sole ration for 21 days between 150 lb.(68 kg) to240 lb.(109 kg) bodyweight. Ractopamine HCl should be fed in a complete swine ration containing at least 16% crude protein.

CAUTION: Pigs fed Ractopamine HCl are at an increased risk for exhibiting the downer pig syndrome (also referred to as “slows,” “subs” or “suspects”). Pig handling methods to reduce the incidence of downer pigs should be thoroughly evaluated prior to initiating use of Ractopamine HCl. Not for use in breeding swine.

MANUFACTURED BY
LINDNER FEED AND MILLING CO. INC.
Comfort, Texas 78013
NET WT. 50 LB (22.68 kg)

LINDNER
677 Pig Grower Finisher LI
MEDICATED

A complete feed for Grower/Finisher Pigs being fed for Exhibition. For control of swine dysentery and the control of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*.

ACTIVE DRUG INGREDIENT

Lincomycin	40 gm/ton
------------------	-----------

GUARANTEED ANALYSIS

Crude Protein – Min.....	17.2%
Lysine—Min.....	0.94%
Crude Fat –Min.....	5.10%
Crude Fiber—Max.....	3.30%
Calcium—Min.....	0.50%
Calcium—Max.....	0.55%
Phosphorous—Min.....	0.45%
Salt—Min.....	0.34%
Salt—Max.....	0.38%
Selenium—Min.....	0.3 ppm
Zinc—Min.....	160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Feed as a sole ration.

Limitations For Use: For Use in animals or on premises with a history of dysentery, but where symptoms have not yet occurred.

CAUTION STATEMENT

Occasionally, swine fed Lincomycin may, within the first 2 days after the onset of treatment, develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5-8 days without discontinuing the Lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feeds containing Lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not to be fed to swine that weigh more than 250 lbs.

MANUFACTURED BY
LINDNER FEED AND MILLING CO. INC.
 Comfort, Texas 78013
 NET WT. 50 LB (22.68 kg)

LINDNER
685 SHOW 16% TY
MEDICATED

A complete feed for increased rate of weight gain and improved efficiency – finisher feeds.

ACTIVE DRUG INGREDIENT

Tylosin.....(10 mg/lb).....	20 grams /ton
-----------------------------	---------------

GUARENTEED ANALYSIS

Crude Protein—Min.....	16.0%
Lysine—Min.....	1.0%
Crude Fat –Min.....	5.0%
Crude Fiber –Max.....	3.0%
Calcium –Min.....	0.6%
Calcium –Max.....	0.7%
Phosphorous –Min.....	0.4%
Salt –Min.....	0.2%
Salt –Max.....	0.3%
Selenium—Min.....	0.3 ppm
Zinc –Min.....	175.0 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotine, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Lindner’s Grower Finisher TY is a complete feed and should be feed continuously to pigs weighing 70 to 220 pounds. Provide fresh water at all times.

CAUTION

Follow feeding directions at all times. This product contains added selenium. Intake of supplemental selenium should not exceed 0.3 parts per million on a complete diet basis or a total of 3 milligrams per head per day. Store in a dry area. **DO NOT** use any product that has become old, molded or insect contaminated.

MANUFACTURED BY

LINDNER FEED AND MILLING CO. INC.
 Comfort, Texas 78013
 NET WT. 50 LB (22.68 kg)

LINDNER
686 SHAPE & GUTS
MEAL
MEDICATED

A Complete Feed for Growing Pigs Being Fed for Exhibition **For control of swine dysentery** associated with *Brachyspira hyodysenteriae* in pigs up to 250 lbs body weight on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.

ACTIVE DRUG INGREDIENT

Bacitratin Methylene Disalicylate	250 g/ton
---	-----------

GUARENTEED ANALYSIS

Crude Protein—Min.....	14.0%
Lysine—Min.....	0.80%
Crude Fat –Min.....	8.2%
Crude Fiber –Max.....	4.3%
Calcium –Min.....	0.4%
Calcium –Max.....	0.5%
Phosphorous –Min.....	0.3%
Salt –Min.....	0.3%
Salt –Max.....	0.4%
Selenium—Min.....	0.3 ppm
Zinc –Min.....	162 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotine, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Lindner’s Shape & Guts is a complete feed for show pigs from 140 lb to show, and should only be fed to pigs that need more cover and body. Feed as the sole ration. Provide fresh water at all times.

CAUTION

Diagnosis of swine dysentery should be confirmed by a veterinarian when results are not satisfactory. This product contains added selenium. Intake of supplemental selenium should not exceed 0.3 parts per million on a complete diet basis or a total of 3 mg. per head per day. Store in a dry area.

DO NOT use any product that has become old, molded or insect contaminated.

MANUFACTURED BY
LINDNER FEED AND MILLING CO. INC.
 Comfort, Texas 78013
 NET WT. 50 LB (22.68 kg)

LINDNER

690

18% Sow Ration

For Lactating Sows and Gilt's

GUARANTEED ANALYSIS

Crude Protein (Min).....	18.10%
Lysine.....	0.81%
Crude Fat (Min).....	3.00%
Crude Fiber (Max).....	3.05%
Calcium (Min).....	0.68%
Calcium (Max).....	0.76%
Phosphorous (Min).....	0.73%
Salt (Min).....	0.33%
Salt (Max).....	0.42%
Selenium (Min).....	0.30ppm
Zinc (min).....	204ppm

INGREDIENT STATEMENT

Grain Products, Plant Protein Products, Fat Products (feed Grade), Calcium Carbonate, Dicalcium Phosphate, Salt, Choline Chloride, Ferrous Sulfate, Manganous Oxide, Zinc Oxide, Vitamin A Acetate in Gelatin, Vitamin B-12 Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Menadione Sodium Bisulfite Complex, Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Folic Acid, d-Biotin, Lysine, Copper Sulfate, Copper Oxide, Ethylenediamine Dihydriodide, Sodium Selenite.

FEEDING DIRECTIONS

Feed 4-5 pounds per day during first 2/3 of gestation period. Increase to 6 pounds for last 1/3 of the gestation period.

CAUTION

Follow feeding directions at all times. This product contains added selenium. Intake of supplemental selenium should not exceed 0.3 parts per million on a complete diet basis or a total of 3 milligrams per head per day.

Store in a dry area.

DO NOT use any product that has become old, molded, or insect contaminated.

MANUFACTURED BY

LINDNER FEED AND MILLING CO., INC.
Comfort, TX 78013

Net weight 50 lbs. (22.68kg)

LINDNER

1000 Sow Base

Base Mix for Mixing Complete Gestation and Lactation Gilt and Sow Feeds

GUARENTEED ANALYSIS

Crude Protein (Min)	11.00%
Lysine (Min)	1.00%
Crude Fat (Min)	0.23%
Crude Fiber (Max)	0.80%
Calcium (Min)	16.00%
Calcium (Min)	17.00%
Phosphorus (Min)	7.18%
Salt (Min)	7.80%
Salt (Max)	8.02%
Selenium (Min)	6.10%
Zinc (Min)	3700 ppm

INGREDIENTS

Plant protein products, dicalcium phosphate, calcium carbonate, salt, Vitamin A supplement, Vitamin D3 supplement, Vitamin E supplement, thiamine, mononitrate, pyridoxine hydrochloride, D-Calcium pantothenate, niacin supplement, folic acid, riboflavin supplement, menadione sodium bisulfite complex, biotin, Choline chloride, zinc sulfate, manganese sulfate, copper sulfate, calcium iodate, ferrous sulfate, sodium selenite, mineral oil, and vitamin B12 supplement.

FEEDING DIRECTIONS

Mix as shown and feed as the sole ration to Gilt's and Sows During Gestation and Lactation Add approved medication as needed. Do not add additional minerals to these rations.

Lactation 17%

100 lb. Sow Base
500 lb. Soybean Meal 47.0%
1400 lb. Ground Corn

Gestation 13%

100 lb. Sow Base
300 lb. Soybean Meal 47.0%
1600 lb. Ground Corn

MANUFACTURED BY

LINDNER FEED AND MILLING CO. INC.
Comfort, Texas 78013

NET WT 50LB (22.68 kg)

LINDNER

1100 BASE 20 -75

Base Mix for Mixing for Growing Swine.

GUARENTEED ANALYSIS

Crude Protein (Min)	30.00%
Lysine (Min)	3.10%
Crude Fat (Min)	3.85%
Crude Fiber (Max)	1.12%
Calcium (Min)	5.28%
Calcium (Min)	5.55%
Phosphorus (Min)	3.08%
Salt (Min)	2.46%
Salt (Max)	2.57%
Selenium (Min)	0.34 ppm
Zinc (Min)	3662 ppm

INGREDIENTS

Grain products, Plant protein products, Dried Whey, Fish Menhaden meal, Bloodmeal Spr dried, Yeast Culture, dicalcium phosphate, calcium carbonate, salt, Vitamin A supplement, Vitamin D3 supplement, Vitamin E supplement, thiamine, mononitrate, pyridoxine hydrochloride, D-Calcium pantothenate, niacin supplement, folic acid, riboflavin supplement, menadione sodium bisulfite complex, biotin, Choline chloride, zinc sulfate, manganese sulfate, copper sulfate, calcium iodate, ferrous sulfate, sodium selenite, mineral oil, and vitamin B12 supplement.

FEEDING DIRECTIONS

Mix as shown and feed as the sole ration to growing pigs. This will make a ration containing approximately 20% crude protein.

200 lb. Base 20-75
500 lb. Soybean Meal 47.0%
1060 lb. Ground Corn
200 lb. Whey
40 lb. Soybean Oil

MANUFACTURED BY

LINDNER FEED AND MILLING CO. INC.
Comfort, Texas 78013

NET WT 50LB (22.68 kg)

LINDNER
1103 LEAN TARGET

Medicated

A Supplement for Finishing Pigs Being Fed for Exhibition

For improved feed efficiency and increased carcass leanness in finishing swine fed a complete ration containing at least 16% crude protein from 150 LB to 240 LB body weight.

ACTIVE DRUG INGREDIENT

Ractopamine hydrochloride (90 mg per LB) ... 180 gr per ton

GUARANTEED ANALYSIS

Crude Protein (min)	20.00%
Lysine (min)	1.50%
Crude Fat (min)	5.00%
Crude Fiber (max).....	6.50%
Calcium (min).....	0.90%
Calcium (max).....	1.13%
Phosphorus (min).....	0.65%
Salt (min).....	0.25%
Salt (max).....	0.75%
Selenium (min).....	0.3ppm
Zinc (min).....	120ppm
Biotin (min).....	1 mg/LB

INGREDIENTS

Ground corn, Soybean Meal, Wheat Middlings, Calcium Carbonate, Animal Fat, BHT (A Perservative), Fish Meal, Yeast Culture, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Sodium Propionate & Ethoxyquin (Preservatives), L-Lysine, Zinc Oxide, Biotin, DL-Methionine, Ferrous Sulfate, Defluorinated Phosphate, d-alpha Tocopherol Acetate (Source of Vitamin E, Mineral Oil, Niacin Supplement, Thiamine Mononitrate, Manganous Oxide, Vitamin E Supplement, Copper Sulfate, Calcium Pantothenate, Magnesium Oxide, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Vitamin A Acetate, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B12 Supplement, Folic Acid, Potassium Iodide, D-Activated Animal Sterol (Source of Vitamin D3), Sodium Selenite.

INSTRUCTIONS FOR FEEDING

Lindner's Lean Target should be top-dressed with a non-medicated pig feed to get a finished complete feed containing at least 16% protein. The feed can only be used to feed pigs weighing from 150 to 240 pounds.

Follow the below mixing instructions.

LEAN	16% PROTEIN	RACTOPAMINE
TARGET	FEED	HYDROCHORIDE LEVEL
0.25 lb	5.00 lb	9 g/ton
0.50 lb	5.00 lb	18 g/ton

WARNING

Ractopamine hydrochloride is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of childrne. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention.

CAUTION: Do not feed to breeding swine.

MANUFACTURED BY
LINDNER FEED & MILLING CO, INC.
Comfort, TX 78013
Net Wt 25 lb (11.33kg)