

**LINDNER  
600 NEO STARTER  
MEDICATED**

A complete feed for starting pigs being fed for exhibition  
For control of swine dysentery associated with *Brachyspira* (formerly  
*Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin.

**ACTIVE DRUG INGREDIENT**

Tiamulin Hydrogen Fumarate ..... 35 g/ton

**GUARANTEED ANALYSIS**

Crude Protein (Min) .....	25.00%
Lysine (Min) .....	1.3%
Crude Fat (Min) .....	5.5%
Crude Fiber (Max) .....	2.0%
Calcium (Min) .....	0.7%
Calcium (Max) .....	1.2%
Phosphorous (Min) .....	0.5%
Salt (Min) .....	0.09%
Salt (Max) .....	0.20%
Selenium (Min) .....	0.4 ppm
Zinc (Min) .....	3,030 ppm

**INGREDIENTS**

Animal Protein Products, Grain Products, Plant Protein Products, Processed Grain By-Products, Animal Fat, Roughage Products, Calcium Carbonate, Ferrous Sulfate, Zinc Sulfate, Selenium Yeast, Vitamin E Supplement, D-Calcium Pantothenate, Vitamin A Supplement, Vitamin D3 Supplement, Riboflavin Supplement, Vitamin B12 Supplement, Manganese Sulfate, Biotin, Copper Sulfate, Niacin Supplement, Dried *Bacillus Licheniformis* Fermentation Product, Dried *Bacillus Subtilis* Fermentation Product, Sodium Silico Aluminate, Dried *Schizosaccharomyces Pombe* Fermentation Product, Sodium Sulfate, Poly-Vinyl Alcohol, Starch, Hydrated Magnesium Silicate, Mineral Oil, D-Alpha-Tocopheryl Acetate, Pyridoxine Hydrochloride, Menadione Dimethylpyrimidinol Bisulfite, Folic Acid, Thiamine Mononitrate, Ethylenediamine Dihydriodide, Zinc Oxide, DL-Methionine, Salt, L-Lysine, Artificial Flavor

**FEEDING DIRECTIONS**

Lindner 600 Neo Starter should be fed continuously as the sole ration for 14 days, and then they should be fed an approved starter ration. Feed continuously as sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following an approved treatment of the disease.

**CAUTION**

Do not feed undiluted. Do not feed to animals other than swine. The effects of Tiamulin on swine reproductive, pregnancy and lactation have not been determined. Use as the only source of Tiamulin. Swine treated with Tiamulin should not have access to feed containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

**WARNING:** Withdraw medicated feed 2 days before slaughter.

**MANUFACTURED BY**  
LINDNER FEED & MILLING CO., INC  
Comfort, TX 78013  
Net Weight 50 lb (22.68 kg)