

**LINDNER  
603 STARTER  
MEDICATED**

A complete feed for starting pigs being fed for exhibition  
For control of swine dysentery associated with *Brachyspira* (formerly  
*Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin.

**ACTIVE DRUG INGREDIENTS**

Tiamulin Hydrogen Fumarate ..... 35 g/ton

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... 24.00%  
Lysine (Min) ..... 1.4%  
Crude Fat (Min) ..... 6.3%  
Crude Fiber (Max) ..... 4.0%  
Calcium (Min) ..... 0.7%  
Calcium (Max) ..... 1.2%  
Phosphorous (Min) ..... 0.6%  
Salt (Min) ..... 0.04%  
Salt (Max) ..... 0.10%  
Selenium (Min) ..... 0.4 ppm  
Zinc (Min) ..... 3,200 ppm

**INGREDIENTS**

Grain Products, Animal Protein Products, Plant Protein Products,  
Animal Fat, Roughage Products, Monocalcium Phosphate, Calcium  
Carbonate, Ferrous Sulfate, Zinc Sulfate, Selenium Yeast, Vitamin E  
Supplement, D-Calcium Pantothenate, Vitamin A Supplement,  
Vitamin D3 Supplement, Riboflavin Supplement, Vitamin B12  
Supplement, Manganese Sulfate, Biotin, Copper Sulfate, Niacin  
Supplement, Dried *Bacillus Licheniformis* Fermentation Product,  
Dried *Bacillus Subtilis* Fermentation Product, Sodium Silico  
Aluminate, Dried *Schizosaccharomyces Pombe* Fermentation  
Product, Sodium Sulfate, Poly-Vinyl Alcohol, Starch, Hydrated  
Magnesium Silicate, Mineral Oil, D-Alpha-Tocopheryl Acetate,  
Pyridoxine Hydrochloride, Menadione Dimethylpyrimidinol  
Bisulfite, Folic Acid, Thiamine Mononitrate, Ethylenediamine  
Dihydriodide, Zinc Oxide, DL-Methionine, L-Lysine, Salt, Artificial  
Flavor

**FEEDING DIRECTIONS**

Pig Starter is a complete feed specifically formulated for pigs fed for  
exhibition. Feed continuously as the sole ration for 14 days, when  
they should be fed an approved grower finisher ration. Feed  
continuously as sole ration on premises with a history of swine  
dysentery but where signs of disease have not yet occurred or  
following an approved treatment of the disease.

**CAUTION**

Do not feed undiluted. Do not feed to animals other than swine. The  
effects of Tiamulin on swine reproductive performance, pregnancy  
and lactation have not been determined. Use as the only source of  
Tiamulin. Swine being treated with Tiamulin should not have access  
to feed containing polyether ionophores (e.g., Monensin, Lasalocid,  
Narasin, Salinomycin and Semduramicin) as adverse reactions may  
occur. If signs of toxicity occur, discontinue use of medicated feeds.

**WARNING:** Withdraw medicated feed 2 days before slaughter.

**MANUFACTURED BY**  
LINDNER FEED & MILLING CO., INC  
Comfort, TX 78013  
Net Weight 50 lb (22.68 kg)