

# LINDNER

## 612 Scud Missile

### MEDICATED

A complete feed for starting pigs being fed for exhibition  
For control of swine dysentery associated with *Brachyspira*  
(formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to  
Tiamulin.

### ACTIVE DRUG INGREDIENTS

Tiamulin Hydrogen Fumarate ..... 35 g/ton

### GUARANTEED ANALYSIS

Crude Protein (Min).....	23.0%
Lysine (Min).....	1.50%
Crude Fat (Min).....	7.10%
Crude Fiber (Max).....	4.70%
Calcium (Min) .....	0.7%
Calcium (Max) .....	1.20%
Phosphorous (Min).....	0.70%
Salt (Min) .....	0.01%
Salt (Max) .....	0.06%
Selenium (Min).....	0.4 ppm
Zinc (Min).....	3,200 ppm

### INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products,  
Animal Fat, Roughage Products 5%, Monocalcium Phosphate,  
Calcium Carbonate, Ferrous Sulfate, Zinc Sulfate, Selenium Yeast,  
Vitamin E Supplement, D-Calcium Pantothenate, Vitamin A  
Supplement, Vitamin D3 Supplement, Riboflavin Supplement,  
Vitamin B12 Supplement, Manganese Sulfate, Biotin, Copper  
Sulfate, Niacin Supplement, Dried *Bacillus Licheniformis*  
Fermentation Product, Dried *Bacillus Subtilis* Fermentation Product,  
Sodium Silico Aluminate, Dried *Schizosaccharomyces Pombe*  
Fermentation Product, Sodium Sulfate, Poly-Vinyl Alcohol, Starch,  
Hydrated Magnesium Silicate, Mineral Oil, D-Alpha-Tocopheryl  
Acetate, Pyridoxine Hydrochloride, Menadione  
Dimethylpyrimidinol Bisulfite, Folic Acid, Thiamine Mononitrate,  
Ethylenediamine Dihydriodide, Zinc Oxide, L-Lysine, DL-  
Methionine, L-Threonine, Artificial Flavors, Salt

### FEEDING DIRECTIONS

Pig Starter Grower is a complete feed specifically formulated for  
pigs fed for exhibition. Feed continuously as the sole ration for 14  
days, when they should be fed an approved grower finisher ration.  
Feed continuously as sole ration on premises with a history of swine  
dysentery but where signs of disease have not yet occurred or  
following an approved treatment of the disease.

### CAUTION

Do not feed undiluted. Do not feed to animals other than swine. The  
effects of Tiamulin on swine reproductive performance, pregnancy  
and lactation have not been determined. Use as the only source of  
Tiamulin. Swine being treated with Tiamulin should not have access  
to feed containing polyether ionophores (e.g., Monensin, Lasalocid,  
Narasin, Salinomycin and Semduramicin) as adverse reaction may  
occur. If signs of toxicity occur, discontinue use of medicated feeds.

**WARNING:** Withdraw medicated feed 2 days before slaughter.

**MANUFACTURED BY**  
LINDNER FEED & MILLING CO., INC  
Comfort, TX 78013  
Net Weight 50 lb (22.68 kg)