

LINDNER
613 Scud Missile
MEDICATED

A complete feed for Finisher pigs being fed for exhibition
For increased rate of weight gain, improved feed efficiency and
increased carcass leanness in finishing swine, weighing not less than
150 lb., fed a complete ration containing at least 16% crude protein
for the last 45 to 90 lb. of gain prior to slaughter (group average)

ACTIVE DRUG INGREDIENTS

Ractopamine Hydrochloride..... 9.0 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min).....	23.00%
Lysine (Min).....	1.50%
Crude Fat (Min).....	7.20%
Crude Fiber (Max).....	4.70%
Calcium (Min).....	0.70%
Calcium (Max).....	0.80%
Phosphorous (Min).....	0.70%
Salt (Min).....	0.01%
Salt (Max).....	0.05%
Selenium (Min).....	0.4 ppm
Zinc (Min).....	3200 ppm

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products,
Animal Fat, Roughage Products 5%, Monocalcium Phosphate,
Calcium Carbonate, Ferrous Sulfate, Zinc Sulfate, Selenium Yeast,
Vitamin E Supplement, D-Calcium Pantothenate, Vitamin A
Supplement, Vitamin D3 Supplement, Riboflavin Supplement,
Vitamin B12 Supplement, Manganese Sulfate, Biotin, Copper
Sulfate, Niacin Supplement, Dried *Bacillus Licheniformis*
Fermentation Product, Dried *Bacillus Subtilis* Fermentation Product,
Sodium Silico Aluminate, Dried *Schizosaccharomyces Pombe*
Fermentation Product, Sodium Sulfate, Poly-Vinyl Alcohol, Starch,
Hydrated Magnesium Silicate, Mineral Oil, D-Alpha-Tocopheryl
Acetate, Pyridoxine Hydrochloride, Menadione Dimethylpyrimidinol
Bisulfite, Folic Acid, Thiamine Mononitrate, Ethylenediamine
Dihydriodide, Zinc Oxide, L-Lysine, DL-Methionine, L-Threonine,
Artificial Flavor, Salt

FEEDING DIRECTIONS

Feed continuously as sole ration to finishing swine weighing 150 lb
(68 kg) to 240 lb (109 kg) for 21 days. Ractopamine HCl should be
fed in a complete swine ration containing at least 16% crude protein
for the last 45 to 90 lb. of gain prior to slaughter (group average).

CAUTION

Pigs fed Ractopamine HCl are at an increased risk for exhibiting the
downer pig syndrome (also referred to as "slows," "subs" or
"suspects"). Pig handling methods to reduce the incidence of downer
pigs should be thoroughly evaluated prior to initiating use of
Ractopamine HCl. Ractopamine may increase the number of injured
and/or fatigued pigs during marketing. Not for use in breeding swine.
NOTE: No increased benefit has been shown when Ractopamine
concentrations in the diet are greater than 4.5 g/ton.

MANUFACTURED BY

LINDNER FEED AND MILLING CO., INC

Comfort, Texas 78013

Net Weight 50 lb (22.68 kg)