

LINDNER
632 Starter / Grower BMD-FBZ
MEDICATED

A complete feed for Starter/Grower pigs

Bacitracin Methylene Disalicylate is indicated for control of swine dysentery associated with *Treponema hyodysenteriae* on the premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.

Fenbendazole is indicated for the removal of: Adult stage Lungworms: *Metastrongylus apri*, *Metastrongylus pudendotectus*. Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); adult stage nodular worms (*Oesophagostomum dentatum*, *Ornithostrongylus quadrispinulatum*); adult stage small stomach worm (*Hyostrongylus rubidus*); Adult and larvae (L2, L3, L4 stages- intestinal mucosal forms) whipworm (*Trichuris suis*); Kidneyworm: Adult and larvae *Stephanurus dentatus*.

ACTIVE DRUG INGREDIENTS

Bacitracin Methylene Disalicylate	250 g/ton
Fenbendazole	29.2 g/ton(14.6 mg/lb)

GUARANTEED ANALYSIS

Crude Protein (Min)	20.0%
Lysine (Min)	1.4%
Crude Fat (Min)	5.0%
Crude Fiber (Max)	3.0%
Calcium (Min)	0.7%
Calcium (Max)	1.2%
Phosphorous (Min)	0.7%
Salt (Min)	0.2%
Salt (Max)	0.7%
Selenium (Min)	0.4 ppm
Zinc (Min)	120 ppm

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Animal Fat, Monocalcium Phosphate, Calcium Carbonate, Sodium Bentonite, Salt, Roughage Products, Dried *Bacillus Licheniformis* Fermentation Product, Dried *Bacillus Subtilis* Fermentation Product, Sodium Aluminosilicate, Ferrous Sulfate, Zinc Sulfate, Choline Chloride, Vitamin E Supplement, D-Calcium Pantothenate, Vitamin A Supplement, Vitamin D3 Supplement, Riboflavin Supplement, Vitamin B12 Supplement, Copper Sulfate, Dried *Schizosaccharomyces Pombe* Fermentation Product, Sodium Sulfate, Poly-Vinyl Alcohol, Starch, Hydrated Magnesium Silicate, Manganese Sulfate, Niacin Supplement, Mineral Oil, D-Alpha-Tocopheryl Acetate, Sodium Selenite, Menadione Sodium Bisulfite Complex, Ethylenediamine Dihydriodide, DL-Methionine, L-Lysine, L-Threonine, Artificial Flavor

FEEDING DIRECTIONS

Feed as the sole ration to provide a total intake of 9mg Fenbendazole/kg of body weight (4.08mg Fenbendazole/lb. of body weight) divided over 3-12 days. Multiply 0.28 times pound of body weight and feed this amount of product over 3-12 days. No withdrawal time needed.

Example:

Body weight of Pig to be Dewormed	Starter/Grower Feed Required per pig	Days Required to Consume Dose
50 lb.	14.0 lb.	7
75 lb.	21.0 lb.	7
100 lb.	28.0 lb.	7

CAUTION: Not for use in swine that weigh more than 250 lb. Under conditions of continued exposure to parasites, retreatment may be needed after 4-6 weeks. Diagnosis of swine dysentery should be confirmed by a veterinarian when results are not satisfactory.

CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT, AND CONTROL OF PARASITISM.

MANUFACTURED BY
LINDNER FEED AND MILLING CO., INC
Comfort, Texas 78013
Net Weight 50 lb (22.68 kg)
NADA # 131-675, Approved by FDA